

TOWER DIGESTIVE HEALTH MEDICAL GROUP
CAREY B. STROM, MD, FASGE
9090 WILSHIRE BOULEVARD, SUITE 101
BEVERLY HILLS, CA 90211
310-550-0400
310-285-0482 (fax)
www.doctorstrom.com

PATIENT INSTRUCTIONS FOR COLONOSCOPY WITH SUPREP PREPARATION

Patient Name: _____

Procedure Date: _____ Arrival Time: _____

LOCATION:

Doheny Endosurgical Ctr.
9090 Wilshire Blvd., Ste. 100
Beverly Hills, CA 90211
310-246-2555

Cedars Sinai Medical Ctr.
GI Lab, South Tower
8700 Gracie Allen Dr.
Los Angeles, CA 90048
310-423-6056

ONE WEEK BEFORE PROCEDURE:

Please do not take any ASPIRIN, MOTRIN, ADVIL, ALEVE, IBUPROFEN (or any other anti-inflammatory medications), TICLID, VITAMIN E, GINKO BILOBA, ST. JOHN'S WORT, VITAMINS CONTAINING IRON, PLAVIX, XARELTO, ELIQUIS, PRADAXA, SAVAYSA, BRILINTA, OR COUMADIN 7 days prior to the procedure. Instructions regarding PLAVIX, XARELTO, ELIQUIS, PRADAXA, SAVAYSA, BRILINTA, OR COUMADIN should be supplied by the physician who is following you on these medications. If you accidentally take one of the above medications, please call the office.

DAY BEFORE PROCEDURE:

1. NO SOLID FOOD ALL DAY BEFORE PROCEDURE, ONLY CLEAR LIQUIDS. You can have water, apple juice, white grape juice, white cranberry juice, black coffee, tea, Gatorade drinks, vegetable/chicken broth, gelatin, popsicles (both no red colored), soft drinks, and hard candy.
2. Follow instructions on the back of this page for mixing, preparing, and taking SUPREP.
3. You may experience discomfort, abdominal distention, abdominal pain, nausea, vomiting, and/or headache.
4. NOTHING BY MOUTH AFTER MIDNIGHT.

OTHER INSTRUCTIONS:

1. You may take high blood pressure and/or heart medications the morning of the procedure with a small amount of water. For insulin, please consult with your primary doctor for instructions.
2. **YOU WILL NOT BE ALLOWED TO DRIVE YOURSELF HOME AFTER THE PROCEDURE. TAXI CABS ARE NOT ALLOWED UNLESS SOMEONE IS ACCOMPANYING YOU. PLEASE MAKE ARRANGEMENTS PRIOR TO THE PROCEDURE.**
3. Confirm your procedure with our office 24-48 hours prior to scheduled procedure.

OVER

SUPREP Bowel Prep Kit is a split-dose (2-day) regimen. Both 6-ounce bottles are required for a complete prep.

FIRST DOSE: Begin Step 1 at **5:00** PM the evening before your procedure and proceed as shown below:

You must complete Steps 1 through 4 using one (1) 6-ounce bottle before going to bed:

STEP 1



Pour **ONE** (1) 6-ounce bottle of SUPREP liquid into the mixing container.

STEP 2



Add cool drinking water to the 16-ounce line on the container and mix.

NOTE: Dilute the solution concentrate as directed prior to use.

STEP 3



Drink **ALL** the liquid in the container.

STEP 4



You **must** drink two (2) more 16-ounce containers of water over the next 1 hour.

SECOND DOSE: Begin Step 1 at **9:00PM**

and proceed as shown above:

For this dose, repeat Steps 1 through 4 shown above using the other 6-ounce bottle.

Scope Out Cancer