

TOWER DIGESTIVE HEALTH MEDICAL GROUP

**CAREY B. STROM, MD, FASGE**

8631 West 3<sup>rd</sup> St Suite 1017E

Los Angeles CA 90048

310-550-0400

310-285-0482 (fax)

[www.doctorstrom.com](http://www.doctorstrom.com)

PATIENT INSTRUCTIONS FOR COLONOSCOPY WITH SUPREP PREPARATION

Patient Name: \_\_\_\_\_

Procedure Date: \_\_\_\_\_ Arrival Time: \_\_\_\_\_

LOCATION:

The Surgery Center of Beverly Hills

8750 Wilshire Boulevard, Suite 150

Beverly Hills CA 90211

(424)-382-1137

**ONE WEEK BEFORE PROCEDURE:**

Please do not take any ASPIRIN, MOTRIN, ADVIL, ALEVE, IBUPROFEN (or any other anti-inflammatory medications), TICLID, VITAMIN E, GINKO BILOBA, ST. JOHN'S WORT, VITAMINS CONTAINING IRON, PLAVIX, XARELTO, ELIQUI S, PRADAXA, SAVAYSA, BRILINTA, OR COUMADIN 7 days prior to the procedure. Instructions regarding PLAVIX, XARELTO, ELIQUIS, PRADAXA, SAVAYSA, BRILINTA, OR COUMADIN should be supplied by the physician who is following you on these medications. If you accidentally take one of the above medications, please call the office.

**DAY BEFORE PROCEDURE:**

1. NO SOLID FOOD ALL DAY BEFORE PROCEDURE, ONLY CLEAR LIQUIDS. You can have water, apple juice, white grape juice, white cranberry juice, black coffee, tea, Gatorade drinks, vegetable/chicken broth, gelatin, popsicles (both no red colored), soft drinks, and hard candy.
2. Follow instructions on the back of this page for mixing, preparing, and taking SUPREP.
3. You may experience discomfort, abdominal distention, abdominal pain, nausea, vomiting, and/or headache.
4. NOTHING BY MOUTH AFTER MIDNIGHT.

**OTHER INSTRUCTIONS:**

1. You may take high blood pressure and/or heart medications the morning of the procedure with a small amount of water. For insulin, please consult with your primary doctor for instructions.
2. **YOU WILL NOT BE ALLOWED TO DRIVE YOURSELF HOME AFTER THE PROCEDURE. TAXI CABS ARE NOT ALLOWED UNLESS SOMEONE IS ACCOMPANYING YOU. PLEASE MAKE ARRANGEMENTS PRIOR TO THE PROCEDURE.**
3. Confirm your procedure with our office 24-48 hours prior to scheduled procedure.

## COVID-19 TESTING:

If a COVID-19 test is needed, Cedars-Sinai offers drive-thru testing. A test should be scheduled 3-5 days prior to your procedure. After talking to our office, please call (310)-423-4674 to schedule an appointment. Thank you!

SUPREP Bowel Prep Kit is a split-dose (2-day) regimen. Both 6-ounce bottles are required for a complete prep.

**FIRST DOSE:** Begin Step 1 at **5:00** PM the evening before your procedure and proceed as shown below:

You must complete Steps 1 through 4 using one (1) 6-ounce bottle before going to bed:

### STEP 1



Pour **ONE (1)** 6-ounce bottle of SUPREP liquid into the mixing container.


### STEP 2



Add cool drinking water to the 16-ounce line on the container and mix.

**NOTE:** Dilute the solution concentrate as directed prior to use.

### STEP 3



Drink **ALL** the liquid in the container.

### STEP 4



You **must** drink two (2) more 16-ounce containers of water over the next 1 hour.

**SECOND DOSE:** Begin Step 1 at **9:00** PM

For this dose, repeat Steps 1 through 4 shown above using the other 6-ounce bottle.

**NOTE:** You **must** finish drinking the final glass of water at least 2 hours, or as directed, before your colonoscopy.

OVER

# Scope Out Cancer